

AD \_\_\_\_\_

Award Number: W81XWH-10-1-0672

TITLE: Molecular Targeted Therapies of Childhood Choroid Plexus Carcinoma

PRINCIPAL INVESTIGATOR: David Malkin

CONTRACTING ORGANIZATION: The Hospital for Sick Children  
Toronto, Ontario Canada M5G 1X8

REPORT DATE: October 2011

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE October 2011		2. REPORT TYPE Annual		3. DATES COVERED 15 September 2010 – 14 September 2011	
4. TITLE AND SUBTITLE  Molecular Targeted Therapies of Childhood Choroid Plexus Carcinoma				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-10-1-0672	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)  David Malkin  E-Mail: David.Malkin@sickkids.ca				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  The Hospital for Sick Children Toronto, Ontario Canada M5G 1X8				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT  None provided.					
15. SUBJECT TERMS None provided.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	9	19b. TELEPHONE NUMBER (include area code)

## Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	1
Key Research Accomplishments.....	1
Reportable Outcomes.....	4
Conclusion.....	4
References.....	4
Appendices.....	5

# Key Research Accomplishments from Dr. Malkin toward CA093469: Molecular Targeted Therapies of Childhood Choroid Plexus Carcinoma (CPC)

## INTRODUCTION

Choroid plexus carcinoma (CPC) is a rare malignant brain tumor originating from the epithelial cells lining the cerebral ventricles. CPC represents less than 0.6% of brain tumors in all age groups, yet is more frequent in children (2-4%), especially in infants under the age of 1, accounting for over 20% of brain tumors in this age group (1). The molecular events that drive the malignant progression of this tumor are not well understood, yet this knowledge is crucial to improve patient survival. Surgical resection combined with neo-adjuvant and/or adjuvant therapy remain the primary methods of treatment for CPC; however tumor progression and relapse is observed in ~70% of cases (2). Despite improvements on the most current treatment protocols, long-term survival of CPC patients remains under 30% and survivors display significant neurocognitive and/or sensory deficits. (2,3). Identifying altered genes that drive the progression of CPC will refine current diagnostic and prognostic classifications of CPC patients, and promote the implementation of targeted therapies to improve patient survival and reduce long-term side effects. The proposed research aims to identify genetic lesions involved in CPC tumorigenesis in order to implement their use as unique markers for diagnostic and prognostic classification of choroid plexus tumor patients, as well as to promote the creation of personalized molecular targeted therapies.

*I hypothesize that recurrent genetic lesions accompanied by a significant change in gene expression in CPC, will be drivers of tumorigenesis in this malignant brain tumor.* Consequently, this research project will seek to answer the following question: What genes may be used as molecular markers for diagnostic and prognostic classification of CPC, and for which tumor-promoting alterations could CPC therapies be created?

## BODY AND KEY RESEARCH ACCOMPLISHMENTS

We have made great progress during the first 11 months of our project; completing all the elements proposed in our Statement of Work (SOW). We have maintained twice monthly teleconferences and data sharing exchanges with the group at St. Jude throughout the last funding period. This proved extremely important, ensuring that we exchange data and ideas in real time, maximizing the efficiency of all project interactions. The following narrative is set in the context of our group SOW, highlighting specifically those tasks assigned to Dr. Malkin. Sections of the original SOW are shown in italics, followed by details of tasks completed.

### SOW Specific Aim 1, Task 1

*Specific Aim 1: To identify candidate drug targets of CPC.*

**Task 1: Generation of additional human and mouse CPC genomic profiles (timeframe: months 1-5).**

*The goal of these studies is to expand our number of genomic profiles (DNA and mRNA arrays) of both human and mouse CPCs to provide a comprehensive dataset with which to identify key candidate oncogenes, tumor suppressor genes (TSGs) and aberrant signal pathways that underlie the development of CPC. NB. Dr. Malkin will continue to generate human genomic profiles and compare these to mouse CPC profiles throughout the duration of the grant.*

**Task 1.d.:** *Identify and classify each of the copy number alterations (CNAs) in initial cohorts of human and mouse tumors as 'frequent' (>20% of tumor samples), moderately frequent (>10% to <20%), or rare (<10%) and according to distribution between human TP53 wild-type or mutant tumors and among the different stages of mouse tumor development. PIs: Gilbertson, Malkin, Guy and Ellison. Timeframe: months 4-5.*

**Task 1.e:** *Accrue and perform central histology review and Affymetrix 6.0 SNP and Exon expression arrays of an additional human CPCs during the course of the granting period. PIs: Malkin, Ellison, Gilbertson. Timeframe: months 1-36 (ongoing throughout grant period).*

***Milestone #1: Completion of initial human and mouse CPC genomic datasets.***

**Work completed toward Specific Aim 1, Task 1 by Dr. Malkin**

**Generation of additional human CPC genomic datasets**

- DNA and RNA extraction was conducted for more than 90 human choroid plexus samples using well-established protocols.
- A variety of quality control tests were run to assess the integrity and quality of all samples. Fifty-nine DNA samples were selected for hybridization to Affymetrix SNP 6.0 GW microarrays, while 40 RNA samples were selected for reverse transcription reactions yielding cDNA for hybridization to Affymetrix Exon 1.0 ST microarrays.
- SNP and Exon microarrays were run at the Toronto Centre for Applied Genomics (TCAG) microarray facility.

This represents a significant contribution in the form of additional new human tumor data toward the analysis of CPCs. These data have been used directly to complete Task 2.

**SOW Specific Aim 1, Task 2**

***Task 2: Genome-wide cataloging of candidate oncogenes and TSGs in human and mouse CPCs (Timeframe: months 5-8).***

*The goal of these studies is to identify candidate oncogenes and TSG that map within the CNAs detected in human and mouse CPCs.*

***Task 2.a: Catalog all genes located within the frequent, moderately frequent and rare CNAs identified in human and mouse CPCs. PIs: Gilbertson and Malkin. Timeframe: months 5-6.***

***Task 2.b: Rank probability of the human and mouse genes within all inferred CNAs as being candidate oncogenes or TSGs of CPC. PIs: Gilbertson and Malkin. Timeframe: months 5-6.***

***Task 2.c: Select the top 30 candidate oncogenes and TSGs of human and mouse CPCs. PIs: Gilbertson, Malkin, Ellison and Guy. Timeframe: month 6.***

***Task 2.d: Real-Time PCR and FISH validation of copy number alterations of our lead candidate oncogenes and TSGs in human and mouse CPCs. PI: Ellison. Timeframe: months 6-8.***

***Milestone #2: Selection of the top candidate oncogenes and tumor suppressor genes of CPC.***

**Work completed toward Specific Aim 1, Task 2 by Dr. Malkin**

**Genomic and transcriptomic analyses to identify candidate tumor suppressor genes and oncogenes**

All raw data generated for each DNA and mRNA tumor microarray was used for copy number and gene expression analysis. These are now all complete and cataloged for alterations as detailed below.

***Copy Number Analysis-Task 2a.***

- Copy number analysis was performed using Partek Genomics Suite (PGS, Partek Inc. St. Louis, MO) and visualized using Integrative Genomic Viewer (IGV, <http://www.broad.mit.edu/igv/>)
- Normalization of all data was then performed with SNP 6.0 files created for the third phase of the International HapMap Project (HapMap3, 1301 control samples)
- We identified all areas of copy number alteration (CNA) that might contain oncogenes and tumor suppressor genes using the Hidden Markov Model (HMM) and Partek's segmentation algorithm. The minimum number of markers per region was set at 10.
- Candidate CNAs were statistically reviewed by a rigorous set of criteria to prioritize regions of gain or loss for further analysis.

***Gene Expression Analysis-Task 2a and b.***

- Gene expression analysis was performed using PGS and Gene Pattern (<http://genepattern.broad.mit.edu/>).
- Four clustering algorithms (principal component analysis (PCA), unsupervised hierarchical clustering (UHC), non-negative matrix factorization (NMF) and Consensus Clustering (CC)), were used to identify tumor subgroups that segregate independently according to their expression profile. In all analyses, a significant segregation was observed between CPC and choroid plexus papilloma (CPP) samples.
- Microarray intensities were analyzed in PGS, using the benign human choroid plexus papilloma (CPP) samples as an expression baseline reference. This analysis highlights expression changes unique to the malignant choroid plexus carcinoma (CPC) phenotype. Significant changes in expression between CPC and CPP were considered as a fold-change of 2 or more. P-value FDR= 0.05 after ANOVA analysis

#### ***Copy number-driven expression-Task 2b and c***

- To analyze copy number-driven expression in choroid plexus tumors, we merged the copy number for each segmented region obtained from copy number analysis to the gene expression values.
- The data was filtered according to direction of copy number alteration and expression changes. Only those regions with copy number gain and increased expression, as well as copy number loss and decreased expression were selected.
- Further data filtering took into consideration the frequency of copy number-driven events in the CPC group compared to the CPP group, as the focus was on CPC-unique events.
- An initial list of recurrent CPC-unique regions with significant copy number-driven expression was compiled. These regions were observed in at least 56% of CPC samples (n=10/18). Seventy seven CNAs containing candidate oncogenes were identified. Most amplifications map to chromosome 1, but also include on chromosomes 5, 7, 8, 9 12, 14, 15, 17, 18, 19, 20, 21, X. (**Appendix 1,2**)

All data were discussed in real time with the entire group to ensure that we prioritized and selected the appropriate lesions for further analysis (See the reports of Drs. Ellison and Gilbertson in particular). Regions prioritized as special interest in cross species analyses as likely to contain oncogenes of mouse and human CPCs were then passed forward to Task 3.

Together with Drs. Gilbertson and Ellison we have been working to validated in human CPCs, a series of alterations that are also observed in mouse tumors. These CNAs that are common to both species are likely to contain critical genes important in the development of these tumors.

#### ***Validating regions with qPCR and FISH-Task 2d.***

- Validation of genomic and transcriptomic findings in human CPC by qPCR and FISH are currently in progress. Preliminary qPCR results have confirmed copy number gain in at least four CPC samples for the following genes: STMN1 (1p36), EXO1 (1q43), STIL (1p32), ZYG11A (1p32), RAD54L (1p32), and SLC1A7 (1p32).
- **NB. Drs. Gilbertson and Ellison have also confirmed CNAs in these same genes in mouse CPCs. These are extremely exciting data since they identify for the first time candidate oncogenes of CPCs that are present in tumors in two separate species. We are now working with Dr. Guy to determine if these genes are targetable with novel therapies.**

#### **SOW Specific Aim 1, Task 3**

***Task 3: Interspecies analysis of gene candidates and pathway alterations common to human and mouse CPCs (Timeframe: months 6-8).***

*The goal of this analysis is to identify a set of common genes that are dysregulated in both human and mouse CPCs. Genes that are non-redundant for CPC tumorigenesis across species are likely to be critical to the disease process.*

**Task 3.a:** Compare lists of orthologs of candidate oncogenes and TSGs between mouse and human CPC. **PIs:** Gilbertson and Malkin. **Timeframe:** months 6-8.

**Milestone #3:** Comprehensive understanding of the molecular similarities between human and mouse CPC.

**Work completed toward Specific Aim 1, Task 3 by Dr. Malkin**

**Comparison between mouse and human data-Task 3a.**

- The list generated from regions exhibiting copy number-driven expression in human CPC was used to compare equivalent regions in mouse CPC. The cross-species analysis was analyzed by the Gilbertson group.

**REPORTABLE OUTCOMES**

**ABSTRACTS:**

1. Merino D, et al. Choroid plexus carcinoma is characterized by unique genomic and expression alterations in human chromosome 1. 102<sup>nd</sup> Annual Meeting of the American Association for Cancer Research, Orlando, FL. April 1-5, 2011
2. Merino D, et al. Aberrant copy number-driven expression in chromosome 1 is involved in choroid plexus development. EMBL Cancer Genomics Workshop, Heidelberg, Germany. October 1-4, 2011.


**CONCLUSIONS**

Our data to date indicates differential patterns of gene amplification and deletion between choroid plexus papilloma and malignant choroid plexus carcinoma. The most striking chromosomal targets are throughout chromosome 1, and these appear to correlate highly with mouse orthologs. This data provides a robust basis on which to pursue functional characterization, cellular localization and drug target discovery studies in the next 2 years of the grant cycle.

**REFERENCES**

1. Aguzzi A., Bradner S., & Paulus W. (2000) Choroid plexus tumors. In: Kleihues P., Cavenee WK., eds. World Health Organization Classification of Tumors: Pathology and Genetics of Tumors of the Nervous System. Lyon, France: IARC Press
3. Lafay-Cousin, L. et al. (2010) Choroid plexus tumors in children less than 36 months: the Canadian Pediatric Brain Tumor Consortium (CPBTC) experience. Childs Nerv Syst 27(2), 259-264
4. Lafay-Cousin, L. et al. (2010) Use of ifosfamide, carboplatin, and etoposide chemotherapy in choroid plexus carcinoma. J Neurosurg Pediatrics 5(6), 615-621

APPENDIX 1 (Figure 1)



**Figure 1:** Chromosome 1 contains 23 significant regions of copy number-driven expression unique to CPC.



## Appendix 2 (Figure 2)

**Figure 2:** Profile of chromosome-wide CNA in CPP and CPC samples. Black squares represent regions of copy number gain ( $\geq 2.75$ ) or loss ( $\leq 1.25$ ) that cover more than 50% of each chromosome arm. P-values calculated by Fisher's exact test (significant p-values in red).

